

# VERAによる長周期変光星のVLBI位置天文観測

中川 亜紀治 (1), 倉山 智春 (2), 面高 俊宏 (1), 半田 利弘 (1)  
星原 一航 (1), 村上 琴音 (1), 岩井 智美 (1), 大山 まで薫 (1)  
VERA プロジェクト (3)

(1) 鹿児島大学 (2) 帝京科学大学 (3) 国立天文台 水沢VLBI観測所  
nakagawa@sci.kagoshima-u.ac.jp



ミラ型変光星を含む長周期変光星 (LPV) は太陽の 1–8 倍の質量を持つ。質量放出は激しく、宇宙の化学組成においても重要な役割をもつ。この種の星では明るさと変光周期の比例関係 (周期光度関係; Period-luminosity relation; PLR) が知られており、VERA プロジェクトの科学目標の一つはこの関係を我々の天の川銀河独自で確立することである。我々はミラ型変光星を中心にこれまで多くの LPV の位置天文観測を行ってきた。ポスターでは VLBA による計測結果とあわせてこれまでの結果を示した。

星周物質の運動の導出法として HIPPARCOS による計測との差分を用いる手法を試みた。この際、バイナリー運動の可能性を検討する必要がある。本解析からはバイナリーについての強い制限を加えることは難しかったが、少なくとも 0.5 km/s/yr より大きな加速度は検出できなかった。近く打ち上げが予定される Nano-JASMINE には GAIA では困難な近傍ミラ型変光星 (D=数百 pc) の固有運動測定が期待される。このポスターで論じる VERA の測定と結びつける方法により、内部運動やバイナリー運動についてより精度の高い議論が期待できる。

## 1. Mira 型変光星の位置天文観測

### 1.1 Source selection and single dish monitoring

In table 1, we show a part of our target sources. Long period variables (LPVs) including Miras and semiregular variables are presented with Pulsation periods (P). Figure 1 shows a period distribution of ~800 Miras in Feast et al. (2000) and our ~80 targets. Period average of ~80 Miras with water maser emission is 407 day (LogP = 2.61), which is longer than that of 338 day (LogP = 2.53) of the sources in Feast et al. (2000).

To determine a beginning of the series VLBI observations, we monitor >250 LPVs at Iriki station (Shintani et al. 2008). Intensity of ~10 Jy is a threshold of a successful detection in VLBI with VERA.




Fig.1. Period distribution of Mira variables in Feast et al. (2000) (white) and our targets (green).

Table1: Target Sources

| Name      | P   | LogP | Type |
|-----------|-----|------|------|
| SY Scl    | 415 | 2.62 | Mira |
| WX Psc    | 660 | 2.82 | Mira |
| RU Ari    | 354 | 2.55 | SR   |
| T Lep     | 368 | 2.57 | Mira |
| BW Cam    | --- | ---  | Mira |
| RW Lep    | 150 | 2.18 | SR   |
| BX Cam    | 454 | 2.66 | Mira |
| U Ori     | 368 | 2.57 | Mira |
| AP Lyn    | 450 | 2.65 | Mira |
| U Lyn     | 434 | 2.64 | Mira |
| GX Mon    | 527 | 2.72 | Mira |
| Z Pup     | 509 | 2.71 | Mira |
| QX Pup    | --- | ---  | Mira |
| R Cnc     | 362 | 2.56 | Mira |
| X Hya     | 301 | 2.48 | Mira |
| R UMa     | 302 | 2.48 | Mira |
| S Crt     | 155 | 2.19 | SR   |
| VX UMa    | 215 | 2.33 | Mira |
| T UMa     | 257 | 2.41 | Mira |
| RS Vir    | 354 | 2.55 | Mira |
| FV Boo    | 340 | 2.53 | SR   |
| W Hya     | 361 | 2.56 | Mira |
| RX Boo    | 278 | 2.44 | SR   |
| Y Lib     | 276 | 2.44 | Mira |
| S CrB     | 360 | 2.56 | Mira |
| SW Lib    | 292 | 2.47 | Mira |
| FS Lib    | 415 | 2.62 | Mira |
| IRC+10374 | --- | ---  | Mira |
| IRC-20540 | 510 | 2.71 | Mira |
| SY Aql    | 356 | 2.55 | Mira |
| SV Peg    | 145 | 2.16 | SR   |
| R Aqr     | 390 | 2.59 | Mira |

### 1.2 VLBI observation

We conduct phase referencing VLBI observation at 22 GHz with typical duration of 1.5 – 2.0 years. Interval is ~1 month.

## 2. VLBIによるLPVの測距と周期光度関係

In the last decade, astrometric VLBI has been played an important role for determining annual parallaxes of the Galactic LPVs. Table 2 shows all parallaxes measured with VLBI method (VLBA and VERA). Red giants and OH/IR stars are also included in the table. Errors of absolute magnitude are based on their distance errors.

In the red-colored plot of figure 2, we show a distribution of the Galactic LPVs on absolute magnitude  $M_K$  –  $\log P$  plane. Red solid line indicate a PLR determined in our previous work (Nakagawa et al. 2014). Result by Ita et al. (2004) is superposed to find sequences for several types of variables.




Fig.2. LPV source distribution in the Galaxy (red) and LMC (gray, Ita et al. 2004) on  $M_K$  –  $\log P$  plane. Distance of 49.89 kpc is assumed to the LMC.

Table2: Results from VLBI astrometry

| Source  | Type | Parallax† [mas] | P [day] | LogP  | $m_K \pm$ [mag] | $M_K$ [mag] | Maser | Ref.       |
|---------|------|-----------------|---------|-------|-----------------|-------------|-------|------------|
| W Hya   | SRa  | 10.18±2.36      | 361     | 2.558 | -3.16           | -8.12±0.51  | OH    | vle03      |
| R Cas   | Mira | 5.67±1.95       | 430     | 2.633 | -1.80           | -8.03±0.78  | OH    | vle03      |
| UX Cyg  | Mira | 0.54±0.06       | 565     | 2.752 | 1.40            | -9.94±0.24  | H2O   | kur05      |
| S CrB   | Mira | 2.39±0.17       | 360     | 2.556 | 0.21            | -7.90±0.15  | OH    | vle07      |
| U Her   | Mira | 3.76±0.27       | 406     | 2.609 | -0.29           | -7.41±0.16  | OH    | vle07      |
| RR Aql  | Mira | 1.58±0.40       | 396     | 2.598 | 0.50            | -8.51±0.56  | OH    | vle07      |
| S Crt   | SRb  | 2.33±0.13       | 155     | 2.190 | 0.786           | -7.38±0.12  | H2O   | nak08      |
| VY CMa  | SRc  | 0.88±0.08       | 956     | 2.980 | -0.72           | -11.00±0.20 | H2O   | cho08      |
| S Per   | SRc  | 0.413±0.017     | 822     | 2.915 | 1.33            | -10.59±0.09 | H2O   | asa10      |
| R Aqr   | Mira | 4.7±0.8         | 390     | 2.591 | -1.02           | -7.66±0.37  | SiO   | kam10      |
| SY Scl  | Mira | 0.75±0.03       | 411     | 2.614 | 2.55            | -8.07±0.09  | H2O   | nyu11      |
| RX Boo  | SRb  | 7.31±0.5        | 162     | 2.210 | -1.96           | -7.64±0.15  | H2O   | kam12      |
| NML Cyg | ---  | 0.620±0.047     | 1280    | 3.107 | 0.791           | -10.25±0.16 | H2O   | zha12      |
| PZ Cas  | SRc  | 0.356±0.026     | 925     | 2.966 | 1.00            | -11.24±0.16 | H2O   | kus13      |
| T Lep   | Mira | 3.06±0.04       | 368     | 2.566 | 0.09            | -7.48±0.03  | H2O   | nak14      |
| RW Lep  | SRa  | 1.62±0.16       | 150     | 2.176 | 0.639           | -8.31±0.22  | H2O   | kam14      |
| R Aqr   | Mira | 4.59±0.24       | 390     | 2.591 | -1.02           | -7.71±0.11  | SiO   | min14      |
| U Lyn   | Mira | 1.27±0.06       | 434     | 2.637 | 1.533           | -7.95±0.10  | H2O   | kam15      |
| R UMa   | Mira | 1.92±0.05       | 296     | 2.471 | 1.19            | -7.39±0.06  | H2O   | This study |

Ref. vle03 : Vlemmings et al. 2003, kur05 : Kurayama et al. 2005, vle07 : Vlemmings & van Langevelde 2007, nak08 : Nakagawa et al. 2008, cho08 : Choi et al. 2008, asa10 : Asaki et al. 2010, kam10 : Kamohara et al. 2010, nyu11 : Nyu et al. 2011, kam12 : Kamezaki et al. 2012, zha12 : Zhang et al. 2012, kus13 : Kusuno et al. 2013, nak14 : Nakagawa et al. 2014, kam14 : Kamezaki et al. 2014, min14 : Min et al. 2014, kam15 : Kamezaki et al. 2015.

## 3. Mira 型変光星 R UMa の観測

### 3.1 Parallax measurement

Using 12 maser spots, an annual parallax of R UMa was determined to be  $1.92 \pm 0.05$  mas, corresponding to a distance of  $520 \pm 14$  mas. Figure 3 shows parallactic motions in R.A. and Dec.

### 3.2 Maser spot distribution

Distribution of the maser spots around R UMa (Figure 4). Color indicate radial velocity. A cross mark is an estimated position of central star from a consideration of the distribution. The map is 146 au square. An estimated shell with a radius of 85 mas is presented with a dotted circle.




Fig.3. R UMa, parallactic motion. (Nakagawa et al. in prep.)




Fig.4. Distribution of water masers in R UMa. A cross mark indicate an estimated star position.

### 3.3 Maser internal motion revealed from VERA and HIPPARCOS astrometry

Kinematics of circumstellar matter from VLBI method based on “pattern matching”.  
—No definitive position reference (photosphere is not observable)  
—We estimate internal motions of maser spots by subtracting “average motion”  
—Difficulty for determining the “average motion” with few masers

In figure 4, we derived the internal motions by subtracting HIPPARCOS proper motion ( $\mu_x, \mu_y$ ) = (-40.51, -22.66) mas/yr from our VLBI measurements of each maser spot. VLBI proper motion accuracy of the spots was ~0.2 mas/yr.

$$\begin{aligned} \mu_{\text{VERA}} &= \mu_{\text{sys}} + \mu_{\text{int}} \\ \rightarrow \mu_{\text{HIP}} &= \mu_{\text{sys}} \\ \hline \mu_{\text{VERA}} - \mu_{\text{HIP}} &= \mu_{\text{int}} \end{aligned}$$

Figure 5 shows an internal motion velocity gradient along DEC axis found from our analysis.  $V_y = 0$  gives a  $Y = 140$  mas, which is indicated with a horizontal line in figure 4 with peripheral gray region indicatig its error.




Fig.5. Internal motion velocity gradient of maser spots along DEC.

Actually, we have to pay attention to effects of “binary system”. The  $\mu_{\text{bin}}$  and  $\mu_{\text{bin}'}$  indicate velocity vectors of binary motion on Hipparcos and VERA observation dates.

$$\begin{aligned} \mu_{\text{VERA}} &= \mu_{\text{sys}} + \mu_{\text{bin}'} + \mu_{\text{int}} \\ \rightarrow \mu_{\text{HIP}} &= \mu_{\text{sys}} + \mu_{\text{bin}} \\ \hline \mu_{\text{VERA}} - \mu_{\text{HIP}} &= \Delta \mu_{\text{bin}} + \mu_{\text{int}} \quad (\Delta \mu_{\text{bin}} = \mu_{\text{bin}'} - \mu_{\text{bin}}) \end{aligned}$$

Time differential of  $\mu_{\text{VERA}} - \mu_{\text{HIP}}$  include an acceleration of binary system. In next section, we consider the effect of binary motion, and try to give a constraint on binary scenario of R UMa.

### 3.4 Constraint on binary scenario

Since we did not detect difference of systematic motions between HIP and VERA larger than ~2 mas/yr (~5 km/s), an acceleration due to a binary motion can be estimated to be < 0.5 km/s/yr (5 km/s was divided by an interval of 10 yr between HIP and VERA). In a radial velocity, we could not find any difference in  $V_{\text{slr}}$  of R UMa from literatures.

Gray scale of figure 6 gives a companion mass in unit of  $M_{\text{sun}}$  as a function of orbit semimajor axis and acceleration. If we assume the acceleration of 0.5 km/s/yr as an upper limit, and also assume a companion mass larger than 0.5  $M_{\text{sun}}$ , a semimajor axis should be larger than ~14 au.




Fig.6. Binary orbit semimajor axis, acceleration of binary motion and companion mass.

Of course, we can not reject a single star scenario.

For very bright stars, like nearby Milas, Nano-JASMINE will be a powerful and promising telescope to determine their proper motions. In near future, more accurate proper motions from new satellites will be tied up with VLBI measurements of maser spots. With the same method as this study, circumstellar dynamics and binary scenario of many sources can be studied.